

REPORTS
IN
INFORMATICS

ISSN 0333-3590

Backbone colorings for networks

Hajo Broersma, Fedor V. Fomin, Petr A.
Golovach, Gerhard J. Woeginger

REPORT NO 247

April 2003



Department of Informatics
UNIVERSITY OF BERGEN
Bergen, Norway

This report has URL <http://www.ii.uib.no/publikasjoner/texrap/ps/2003-247.ps>

Reports in Informatics from Department of Informatics, University of Bergen, Norway, is available at <http://www.ii.uib.no/publikasjoner/texrap/>.

Requests for paper copies of this report can be sent to:

Department of Informatics, University of Bergen, Høyteknologisenteret,
P.O. Box 7800, N-5020 Bergen, Norway

Backbone colorings for networks ^{*}

Hajo Broersma [†] Fedor V. Fomin [‡] Petr A. Golovach [§]
Gerhard J. Woeginger [¶]

Abstract

We introduce and study backbone colorings, a variation on classical vertex colorings: Given a graph $G = (V, E)$ and a spanning subgraph H of G (the backbone of G), a backbone coloring for G and H is a proper vertex coloring $V \rightarrow \{1, 2, \dots\}$ of G in which the colors assigned to adjacent vertices in H differ by at least two. We study the cases where the backbone is either a spanning tree or a spanning path.

We show that for tree backbones of G the number of colors needed for a backbone coloring of G can roughly differ by a multiplicative factor of at most 2 from the chromatic number $\chi(G)$; for path backbones this factor is roughly $\frac{3}{2}$. For the special case that G is a split graph the difference from $\chi(G)$ is at most an additive constant 2 or 1, for tree backbones and path backbones, respectively. We show that the computational complexity of the problem ‘Given a graph G , a spanning tree T of G , and an integer ℓ , is there a backbone coloring for G and T with at most ℓ colors?’ jumps from polynomial to NP-complete between $\ell = 4$ (easy for all spanning trees) and $\ell = 5$ (difficult even for spanning paths).

1 Introduction and related research

The work presented here is motivated by the general framework for coloring problems related to frequency assignment. In this application area graphs are used to model the topology and mutual interference between transmitters (receivers, base stations): the vertices of the graph represent the transmitters; two vertices are adjacent in the graph if the corresponding transmitters are so close (or so strong) that they are likely to interfere if they broadcast on the same or ‘similar’ frequency channels. The problem in practice is to assign the frequency channels to the transmitters in such a way that interference is kept at an ‘acceptable level’. This has led to various different types of coloring problems in graphs, depending on different ways to model the level of interference, the notion of similar frequency channels, and the definition of acceptable level of interference (See e.g. [13],[18]). One way of putting these problems into a more general framework is the following.

^{*}The work of HJB, FVF and PAG is sponsored by NWO-grant 047.008.006. The work of FVF is also sponsored by EC contract IST-1999-14186: Project ALCOM-FT (Algorithms and Complexity - Future Technologies). Part of the work was done while FVF and PAG were visiting the University of Twente.

[†]broersma@math.utwente.nl. Faculty of Mathematical Sciences, University of Twente, 7500 AE Enschede, The Netherlands.

[‡]fomin@ii.uib.no. Department of Informatics, University of Bergen, N-5020 Bergen, Norway.

[§]golovach@ssu.komi.com. Faculty of Mathematics, Syktyvkar State University, 167001 Syktyvkar, Russia.

[¶]g.j.woeginger@math.utwente.nl. Faculty of Mathematical Sciences, University of Twente, 7500 AE Enschede, The Netherlands.

Given two graphs G_1 and G_2 with the property that G_1 is a spanning subgraph of G_2 , one considers the following type of coloring problems: Determine a coloring of (G_1 and) G_2 that satisfies certain restrictions of type 1 in G_1 , and restrictions of type 2 in G_2 .

Many known coloring problems related to frequency assignment fit into this general framework. We mention some of them here explicitly.

First of all suppose that $G_2 = G_1^2$, i.e. G_2 is obtained from G_1 by adding edges between all pairs of vertices that are at distance 2 in G_1 . If one just asks for a proper vertex coloring of G_2 (and G_1), this is known as the distance-2 coloring problem. Much of the research has been concentrated on the case that G_1 is a planar graph. We refer to [1], [3], [4], [16], [19], and [20] for more details. In some versions of this problem one puts the additional restriction on G_1 that the colors should be sufficiently separated, in order to model practical frequency assignment problems in which interference should be kept at an acceptable level. One way to model this is to use positive integers for the colors (modeling certain frequency channels) and to ask for a coloring of G_1 and G_2 such that the colors on adjacent vertices in G_2 are different, whereas they differ by at least 2 on adjacent vertices in G_1 . This problem is known as the radio coloring problem and has been studied (under various names) in [2], [5], [6], [7], [8], [9], and [17].

The so-called radio labeling problem models a practical setting in which all assigned frequency channels should be distinct, with the additional restriction that adjacent transmitters should use sufficiently separated frequency channels. Within the above framework this can be modeled by considering the graph G_1 that models the adjacencies of n transmitters, and taking $G_2 = K_n$, the complete graph on n vertices. The restrictions are clear: one asks for a proper vertex coloring of G_2 such that adjacent vertices in G_1 receive colors that differ by at least 2. We refer to [12] and [15] for more particulars.

In this paper, we model the situation that the transmitters form a network in which a certain substructure of adjacent transmitters (called the backbone) is more crucial for the communication than the rest of the network. This means we should put more restrictions on the assignment of frequency channels along the backbone than on the assignment of frequency channels to other adjacent transmitters. The backbone could e.g. model so-called hot spots in the network where a very busy pattern of communications takes place, whereas the other adjacent transmitters supply a more moderate service. We consider the problem of coloring the graph G_2 (that models the whole network) with a proper vertex coloring such that the colors on adjacent vertices in G_1 (that model the backbone) differ by at least 2. Throughout the paper we consider two types of backbones: spanning trees and a special type of spanning trees also known as Hamiltonian paths.

1.1 Terminology and notation

All graphs considered in this paper are assumed to be connected. Let $G = (V, E)$ be a connected finite undirected simple graph, and let $T = (V, E_T)$ be a spanning tree of G . A vertex coloring $f : V \rightarrow \{1, 2, 3, \dots\}$ of V is *proper*, if $|f(u) - f(v)| \geq 1$ holds for all edges $uv \in E$. A vertex coloring is a *backbone* coloring for (G, T) , if it is proper and if additionally $|f(u) - f(v)| \geq 2$ holds for all edges $uv \in E_T$ in the spanning tree T . The chromatic number $\chi(G)$ is the smallest integer k for which there exists a proper coloring $f : V \rightarrow \{1, \dots, k\}$. The backbone coloring number $\text{BBC}(G, T)$ of (G, T) is the smallest integer ℓ for which there exists a backbone coloring $f : V \rightarrow \{1, \dots, \ell\}$. When dealing with colorings, we say that two colors z_1 and z_2 are *adjacent* if and only if $|z_1 - z_2| = 1$.

A *Hamiltonian path* of the graph $G = (V, E)$ is a path containing all vertices of G , i.e. a sequence (v_1, v_2, \dots, v_n) such that $V = \{v_1, v_2, \dots, v_n\}$, all v_i are distinct, and $v_i v_{i+1} \in E$ for all $i = 1, 2, \dots, n - 1$. A *split graph* is a graph whose vertex set can be partitioned into a *clique* (i.e. a set of mutually adjacent vertices) and an *independent set* (i.e. a set of mutually nonadjacent vertices), with possibly edges in between. The size of a largest clique in G is denoted by $\omega(G)$. Split graphs are perfect graphs, and hence satisfy $\chi(G) = \omega(G)$.

1.2 Results

We start our investigations of the backbone coloring number by analyzing its relation to the classical chromatic number. How far away from $\chi(G)$ can $\text{BBC}(G, T)$ be in the worst case? To answer this question, we introduce for integers $k \geq 1$ the values

$$\mathcal{T}(k) := \max \{ \text{BBC}(G, T) : G \text{ a graph with spanning tree } T, \text{ and } \chi(G) = k \} \quad (1)$$

It turns out that this function $\mathcal{T}(k)$ behaves quite primitively:

Theorem 1 $\mathcal{T}(k) = 2k - 1$ for all $k \geq 1$.

The upper bound $\mathcal{T}(k) \leq 2k - 1$ in this theorem in fact is straightforward to see. Indeed, consider a proper coloring of G with colors $1, \dots, \chi(G)$, and replace every color i by a new color $2i - 1$. The resulting coloring uses only odd colors, and hence constitutes a ‘universal’ backbone coloring for any spanning tree T of G . The proof of the matching lower bound $\mathcal{T}(k) \geq 2k - 1$ is more involved and will be presented in Section 2.

Next, let us discuss the situation where the backbone tree is a Hamiltonian path. Similarly as in (1), we introduce for integers $k \geq 1$ the values

$$\mathcal{P}(k) := \max \{ \text{BBC}(G, P) : G \text{ a graph with Hamiltonian path } P, \text{ and } \chi(G) = k \} \quad (2)$$

In Section 3 we will exactly determine all these values $\mathcal{P}(k)$ and observe that they roughly grow like $3k/2$. Their precise behavior is summarized in the following theorem.

Theorem 2 For $k \geq 1$ the function $\mathcal{P}(k)$ takes the following values:

- (a) For $1 \leq k \leq 4$: $\mathcal{P}(k) = 2k - 1$;
- (b) $\mathcal{P}(5) = 8$ and $\mathcal{P}(6) = 10$;
- (c) For $k \geq 7$ and $k = 4t$: $\mathcal{P}(4t) = 6t$;
- (d) For $k \geq 7$ and $k = 4t + 1$: $\mathcal{P}(4t + 1) = 6t + 1$;
- (e) For $k \geq 7$ and $k = 4t + 2$: $\mathcal{P}(4t + 2) = 6t + 3$;
- (f) For $k \geq 7$ and $k = 4t + 3$: $\mathcal{P}(4t + 3) = 6t + 5$;

Next, we discuss the special case of backbone colorings on split graphs. Split graphs were introduced by Hammer & Földes [14]; see also the book [11] by Golumbic. They form an interesting subclass of the class of perfect graphs. The combinatorics of most graph problems becomes easier when the problem is restricted to split graphs. The following theorem is a strengthening of Theorems 1 and 2 for the special case of split graphs.

Theorem 3 Let $G = (V, E)$ be a split graph.

- (a) For every spanning tree T in G , $\text{BBC}(G, T) \leq \chi(G) + 2$.
- (b) If $\omega(G) \neq 3$, then for every Hamiltonian path P in G , $\text{BBC}(G, P) \leq \chi(G) + 1$.

Both bounds are tight.




Figure 1: A split graph G with a Hamiltonian path P (bold edges), such that $\chi(G) = 3$ and $\text{BBC}(G, P) = 5$.

Let us show here why for split graphs with clique number 3 the statement in Theorem 3.(b) does not work. Consider the split graph G on five vertices from Figure 1. Vertices v, u, w form a clique, vertex a is adjacent to v, u and vertex b is adjacent to u, w . The clique number (and hence the chromatic number) of this graph is equal to 3. Let $P = (a, v, u, w, b)$ be the Hamiltonian path. We claim that $\text{BBC}(G, P) > \chi(G) + 1$. To the contrary, assume that G, P has a backbone coloring with colors 1,2,3,4. It is easy to see that u can not be colored with color 2 or 3; otherwise we are forced to use the same color for v and w , a clear contradiction. Now suppose that u is colored with color 1 (the case when u is colored with color 4 is similar). Then one of its neighbors in P must have color 3 and the other one color 4. Without loss of generality assume that v has color 3. Vertex a is adjacent in P to v , so the colors 2,3,4 are forbidden for a and the only valid color for a is 1. But a is adjacent in G to u which has color 1 as well. This contradiction completes the proof of the claim.

Finally, we discuss the computational complexity of computing the backbone coloring number: “Given a graph G , a spanning tree T , and an integer ℓ , is $\text{BBC}(G, T) \leq \ell$?” Of course, this general problem is NP-complete. It turns out that for this problem the complexity jump occurs between $\ell = 4$ (easy for all spanning trees) and $\ell = 5$ (difficult even for Hamiltonian paths).

Theorem 4

- (a) The following problem is polynomially solvable for any $\ell \leq 4$: Given a graph G and a spanning tree T , decide whether $\text{BBC}(G, T) \leq \ell$.
- (b) The following problem is NP-complete for all $\ell \geq 5$: Given a graph G and a Hamiltonian path P , decide whether $\text{BBC}(G, P) \leq \ell$.

2 Tree backbones and the chromatic number

This section is devoted to a proof of the lower bound statement $T(k) \geq 2k - 1$ in Theorem 1. Consider some arbitrary $k \geq 1$. We will construct a graph G with

chromatic number $\chi(G) = k$, and a spanning tree T of G , such that $\text{BBC}(G, T) = 2k - 1$.

The graph G is a complete k -partite graph that consists of k independent sets V_1, \dots, V_k that are all of cardinality k^k . Clearly, $\chi(G) = k$. The spanning tree T is defined as the final tree in the following inductive construction: The tree T_0 is a star with root in V_1 and $k - 1$ leaves in the $k - 1$ sets V_2, \dots, V_k , one in each set. For $j = 1, \dots, k$ the tree T_j is constructed from the tree T_{j-1} , by creating $k - 1$ new vertices for every vertex v in T_{j-1} and by attaching them to v . If v is in the set V_q , then every independent set V_i with $i \neq q$ contains exactly one of these new vertices. Note that all newly created vertices are leaves in the tree T_j . It is easy to see that the tree T_j consists of k^{j+1} vertices that are equally distributed among the sets V_1, \dots, V_k . We denote the vertex set of T_j by $V(T_j)$. Note that $V(T_j) \subset V(T_{j+1})$.

Consider a backbone coloring of (G, T) with ℓ colors where $T = T_k$ is the final tree in the above sequence of trees. Since G is complete k -partite, any color that is used in some set V_i cannot be used in any V_j with $j \neq i$. We denote by C_i the set of colors that are used on vertices in V_i . We now go through a number of steps; in every step, the colors in one of the color sets C_i are labeled with the labels A and B .

(Step s). If there exists some (yet unlabeled) color set C_i such that $|C_i| - 1$ of the colors in C_i are adjacent to a color with label A , then: Label these $|C_i| - 1$ colors with label B . Label the remaining color in C_i with label A .

Eventually, there will be no more color class that satisfies the condition in the if-part: Either, all colors have been labeled, or each of the remaining unlabeled color classes contains at least two colors that are not adjacent to any color with label A . If this is the case at the start, then $|C_i| \geq 2$ for all i , and we obtain $\ell \geq 2k$. We denote by $a \leq k$ the number of steps performed, and may assume $a \geq 1$. We denote by $\pi(s)$ ($s = 1, \dots, a$) the index of the color set that is labeled in step s . Moreover, we denote by $c_{\pi(s)}$ the unique color in $C_{\pi(s)}$ that is labeled A .

Lemma 5 *Let s be an integer with $1 \leq s \leq a$. Then the following statements hold.*

(L1) *In the backbone coloring, all vertices v in $V(T_{k-s}) \cap V_{\pi(s)}$ are colored by color $c_{\pi(s)}$.*

(L2) *The color $c_{\pi(s)}$ is not adjacent to any color $c_{\pi(q)}$ with $q < s$.*

Proof. The proofs of (L1) and (L2) are done simultaneously by induction on s . In step $s = 1$, only a color class $C_{\pi(1)}$ with $|C_{\pi(1)}| = 1$ can be labeled. Then the (unique) color in $C_{\pi(1)}$ is labeled by A , and thus becomes color $c_{\pi(1)}$. But by the definition of $C_{\pi(1)}$, in this case *all* vertices in $V_{\pi(1)}$ are colored by $c_{\pi(1)}$. Statement (b) is trivial for $s = 1$.

Now assume that we have proved the statements up to step $s - 1 < a$, and consider step s . Every color in $C_{\pi(s)} - \{c_{\pi(s)}\}$ (if any) is labeled by B , and is adjacent to some color that has been labeled by A in an earlier step. Let D be the set of these adjacent colors. By the inductive assumption, the colors in D are the only possible colors (from their corresponding color sets) that can be used on the vertices in $V(T_{k-s+1})$. Every vertex v in $V(T_{k-s}) \cap V_{\pi(s)}$ is adjacent to $k - 1$ leaves in T_{k-s+1} , and therefore all the colors in D show up on these leaves. Consequently, they block all colors from $C_{\pi(s)}$ for vertex v except color $c_{\pi(s)}$. This proves statement (L1). In case color $c_{\pi(s)}$ was adjacent to some color x labeled by A in an earlier step, the above argument with $D \cup \{x\}$ instead of D yields that there is no possible color for vertex v . This proves statement (L2). ■

Let L^A denote the set of colors that are labeled by A . Since every step labels exactly one color by A , $|L^A| = a$. Let L^+ denote the set of colors z for which $z - 1$ is in L^A ; clearly, $|L^+| \geq |L^A| - 1 = a - 1$. By statement (L2) in Lemma 5, the sets L^+ and L^A are disjoint. Moreover, there are $k - a$ color sets with unlabeled colors. Since they do not meet the condition in the if-part of the labeling step, each of them contains at least two colors that are not adjacent to any color with label A . These $2(k - a)$ colors are not contained in $L^A \cup L^+$. To summarize, we have found $|L^A| + |L^+| + 2(k - a)$ pairwise distinct colors in the range $1, \dots, \ell$. Therefore,

$$\ell \geq |L^A| + |L^+| + 2(k - a) \geq a + (a - 1) + 2(k - a) = 2k - 1.$$

Note that these arguments also go through in the extremal case $a = k$. This completes the proof of the lower bound statement in Theorem 1.

3 Path backbones and the chromatic number

This section is devoted to a proof of Theorem 2. The upper bound is proved in Section 3.1 by case distinctions. The lower bound is proved in Section 3.2; this proof uses a similar idea as the proof in Section 2, but the actual arguments are quite different.

3.1 Proof of the upper bounds

We start with statement (c) in Theorem 2. Hence, consider a graph $G = (V, E)$ with $\chi(G) = 4t$ for some $t \geq 2$, and let V_1, \dots, V_{4t} denote the corresponding independent sets in the $4t$ -coloring. Furthermore, let $P = (V, E_P)$ be a Hamiltonian path in G . Consider the following color sets:

- For $i = 1, \dots, 3t$, we define the color set $C_i = \{2i - 1\}$.
- For $i = 1, \dots, t$, we define the color set $C'_i = \{2i, 2t + 2i, 4t + 2i\}$.

Note that these $4t$ color sets are pairwise disjoint, and that all the used colors are from the range $1, \dots, 6t$.

We construct a backbone coloring for (G, P) that for $i = 1, \dots, 3t$ colors the vertices in the independent set V_i with the color in color set C_i , and that for $i = 1, \dots, t$ colors the vertices in the independent set V_{3t+i} with one of the three colors in color set C'_i . The vertices in V_{3t+1}, \dots, V_{4t} are colored greedily and in arbitrary order: Consider some vertex v in V_{3t+i} that is to be colored with one of the colors $2i, 2t + 2i, 4t + 2i$. In the worst case, the neighbors of v along the Hamiltonian path P have already been colored by colors x and y , and thus forbid the four colors $x - 1, x + 1, y - 1, y + 1$ for vertex v . Since $t \geq 2$, the three colors in $C'_i = \{2i, 2t + 2i, 4t + 2i\}$ are pairwise at distance at least four, whereas $x - 1, x + 1$ and $y - 1, y + 1$ are at distance two. Therefore, the intersection $C'_i \cap \{x - 1, x + 1, y - 1, y + 1\}$ contains at most two elements, and C'_i contains at least one feasible color for vertex v . This completes the proof of $\mathcal{P}(4t) \leq 6t$ for all $t \geq 2$.

The cases $k = 4t + 1$, $k = 4t + 2$, $k = 4t + 3$ with $t \geq 2$ follow by simple modifications of the above argument: For $k = 4t + 1$, we add the color set $C_{3t+1} = \{6t + 1\}$. For $k = 4t + 2$, we furthermore add the color set $C_{3t+2} = \{6t + 3\}$. And for $k = 4t + 3$, we furthermore add the color set $C_{3t+3} = \{6t + 5\}$. This proves $\mathcal{P}(4t + 1) \leq 6t + 1$, $\mathcal{P}(4t + 2) \leq 6t + 3$, and $\mathcal{P}(4t + 3) \leq 6t + 5$ for all $t \geq 2$, and settles the upper bounds in Theorem 2 for all $k \geq 8$.

The upper bounds in Theorem 2 for all $k \leq 4$ follow trivially from Theorem 1. For $k = 5$, we use the above argument with five color sets

$$D_1 = \{1\}, \quad D_2 = \{3\}, \quad D_3 = \{5\}, \quad D_4 = \{8\}, \quad D_5 = \{2, 6, 7\}.$$

For $k = 6$, we add a sixth color set $D_6 = \{10\}$. Finally, for $k = 7$ we use the seven color sets

$$D'_1 = \{1\}, \quad D'_2 = \{3\}, \quad D'_3 = \{5\}, \quad D'_4 = \{7\}, \quad D'_5 = \{9\}, \quad D'_6 = \{11\}, \quad D'_7 = \{2, 6, 10\}.$$

These three constructions prove $\mathcal{P}(5) \leq 8$, $\mathcal{P}(6) \leq 10$, and $\mathcal{P}(7) \leq 11$. The proof of the upper bounds in Theorem 2 is complete.

3.2 Proof of the lower bounds

We consider a complete k -partite graph G with $k \geq 2$ that consists of k independent sets V_1, \dots, V_k that are all of cardinality $2\Pi_k$. Here Π_k denotes the number of different permutations of $1, 1, 2, 2, 3, 3, \dots, k, k$ in which no two subsequent symbols are the same. It is routine to deduce by inclusion-exclusion that $\Pi_k = \sum_{j=0}^k (-1)^j \binom{k}{j} \frac{(2k-j)!}{2^{k-j}}$. The Hamiltonian path P consists of Π_k segments with $2k$ vertices each. Every such segment visits every independent set exactly twice, since we let each segment correspond to one permutation π of the $2k$ indices $1, 1, 2, 2, 3, 3, \dots, k, k$ that contributes to the total number of Π_k defined before, and we let the segment visit the independent sets exactly in the order $V_{\pi(1)}, V_{\pi(2)}, \dots, V_{\pi(2k)}$. Since G is complete k -partite it is clear that these segments can be combined (in many ways) to form a Hamiltonian path in G . It is also obvious that $\chi(G) = k$.

Consider some fixed backbone coloring of (G, P) with ℓ colors. Since G is complete k -partite, any color that shows up in some set V_i cannot show up in any V_j with $j \neq i$. We denote by C_i the set of colors that are used on vertices in V_i . If $|C_i| = 1$, then V_i is called *mono-chromatic*; if $|C_i| = 2$, then V_i is *bi-chromatic*; if $|C_i| \geq 3$, then V_i is *poly-chromatic*. We denote by s_1 , s_2 , and s_3 the number of mono-chromatic, bi-chromatic, and poly-chromatic sets, respectively. Then clearly

$$s_1 + s_2 + s_3 = k \tag{3}$$

and

$$s_1 + 2s_2 + 3s_3 \leq \ell. \tag{4}$$

Colors that are used on mono-chromatic, bi-chromatic, poly-chromatic sets, are called mono-chromatic, bi-chromatic, poly-chromatic colors, respectively. We say that two bi-chromatic colors x, y with $1 \leq x < y \leq \ell$ are *partner* colors, if $C_i = \{x, y\}$ holds for some bi-chromatic set V_i .

Clearly, we may assume there are mono-chromatic colors. Now consider the following process that labels some of the colors in $\{1, 2, \dots, \ell\}$ with the labels A and B , and that creates a number of arcs among the labeled colors.

(Phase 1). All mono-chromatic colors are labeled by label A .

(Phase 2). Repeat the following step over and over again, as long as the condition in the if-part is met:

If there exists an unlabeled bi-chromatic color y that is adjacent to another color z that has already been labeled A at an earlier point in time, then y is labeled B and its partner color x is labeled A . Moreover, we create an arc going from z to y , and another arc going from y to x .

This process eventually terminates, since the step in the second phase can be performed at most s_2 times. We denote by a and b the number of A -labels and B -labels in the final situation after termination.

Lemma 6 *After termination, the following properties are satisfied.*

(T1) $a = b + s_1$.

(T2) *For every labeled color z , there is a unique directed path from some mono-chromatic color to z .*

(T3) *For two adjacent colors z and $z + 1$, at least one of them is not labeled A .*

Proof. Proof of (T1). After the first phase, there are exactly s_1 colors with A -labels and no vertices with B -labels. Every time the step in the second phase is performed, exactly one new label A and one new label B are created.

Proof of (T2). This is straightforward from the definition of the second phase.

Proof of (T3). Suppose for the sake of contradiction that the adjacent colors z and $z + 1$ are both labeled A . By (T2), there exists a directed path from some mono-chromatic color $x_{\phi(0)}$ to z (note that $x_{\phi(0)} = z$ might hold). This path goes through colors $x_{\phi(0)}, y_{\phi(1)}, x_{\phi(1)}, y_{\phi(2)}, x_{\phi(2)}, \dots, y_{\phi(f)}, x_{\phi(f)}$, with $x_{\phi(f)} = z$. Every color $x_{\phi(i)}$ has an A -label, and every color $y_{\phi(i)}$ has a B -label. Every color $y_{\phi(i)}$ is adjacent to color $x_{\phi(i-1)}$. Moreover, the colors $x_{\phi(i)}$ and $y_{\phi(i)}$ are used on the independent set $V_{\phi(i)}$. By similar considerations, we find a directed path from some mono-chromatic color $x_{\psi(0)}$ to $z + 1$ that goes through colors $x_{\psi(0)}, y_{\psi(1)}, x_{\psi(1)}, \dots, y_{\psi(g)}, x_{\psi(g)}$, with $x_{\psi(g)} = z + 1$. Every color $x_{\psi(i)}$ has an A -label, and every color $y_{\psi(i)}$ has a B -label. Colors $x_{\psi(i)}$ and $y_{\psi(i)}$ are used on the independent set $V_{\psi(i)}$.

Note that the colors in the directed path from $x_{\phi(0)}$ to z are pairwise distinct, and that the colors in the directed path from $x_{\psi(0)}$ to $z + 1$ are pairwise distinct. By the construction of the complete k -partite graph G , there exists a subpath Q of the Hamiltonian path P that visits the independent sets in the ordering

$$V_{\phi(0)}, V_{\phi(1)}, V_{\phi(2)}, \dots, V_{\phi(f)}, V_{\psi(g)}, V_{\psi(g-1)}, V_{\psi(g-2)}, \dots, V_{\psi(1)}, V_{\psi(0)}.$$

Let $v_{\phi(i)}$ and $v'_{\psi(j)}$ be the corresponding vertices on Q . What are the possible colors for these vertices in the backbone coloring under investigation? Vertex $v_{\phi(0)}$ is in a mono-chromatic set, and so it must get color $x_{\phi(0)}$. Vertex $v_{\phi(1)}$ is in a bi-chromatic set, and can be colored by color $x_{\phi(1)}$ or by color $y_{\phi(1)}$. However, $v_{\phi(0)}$ is adjacent to $v_{\phi(1)}$, and its color $x_{\phi(0)}$ is adjacent to $y_{\phi(1)}$. Therefore, $v_{\phi(1)}$ must be colored by $x_{\phi(1)}$. Analogous arguments show that every vertex $v_{\phi(i)}$ is colored by color $x_{\phi(i)}$, and that every vertex $v'_{\psi(i)}$ is colored by color $x_{\psi(i)}$.

Now we arrive at the desired contradiction: Vertex $v_{\phi(f)}$ is colored by color $x_{\phi(f)} = z$, vertex $v'_{\psi(g)}$ is colored by color $x_{\psi(g)} = z + 1$, and hence two adjacent vertices on the backbone are colored by adjacent colors. ■

Let L denote the set of colors z for which $z + 1$ is labeled A after termination. If color 1 is labeled A , then $|L| = a - 1$, and otherwise $|L| = a$. In any case, $|L| \geq a - 1$. No color in L can be labeled A , since this would contradict property (T3) in Lemma 6. At most b of the colors in L can be labeled B . Hence, L contains at least $a - 1 - b = s_1 - 1$ unlabeled colors, where the equation follows from (T1). None of these $s_1 - 1$ unlabeled colors can be bi-chromatic; otherwise, there would be another possible step in the second phase. Hence, these $s_1 - 1$ unlabeled colors in L must all be poly-chromatic. Among the ℓ colors used by the backbone coloring, there are s_1 mono-chromatic ones, $2s_2$ bi-chromatic ones, and at least $s_1 - 1$ poly-chromatic ones. Therefore,

$$2s_1 + 2s_2 - 1 \leq \ell. \quad (5)$$

Adding inequality (4) to inequality (5), and subtracting three times the equation in (3) yields

$$3k + s_2 - 1 \leq 2\ell. \quad (6)$$

Since s_2 is non-negative, (6) implies that $\ell \geq \lceil (3k - 1)/2 \rceil$. For the three cases (c) $k = 4t$, (d) $k = 4t + 1$, (e) $k = 4t + 2$ in Theorem 2 this already implies the claimed lower bounds (c) $\ell \geq 6t$, (d) $\ell \geq 6t + 1$, and (e) $\ell \geq 6t + 3$, respectively. The case (f) $k = 4t + 3$ can be handled as follows: If $s_1 + s_2 \geq 3t + 3$, then (5) implies $\ell \geq 6t + 5$. If $s_1 + s_2 \leq 3t + 2$, then subtracting three times (3) from (4) yields

$$\ell - 3k \geq -2s_1 - s_2 \geq -2(s_1 + s_2) \geq -6t - 4,$$

and hence $\ell \geq 6t + 5$ as desired in statement (f).

It remains to prove the ‘small’ cases $k \leq 6$ in statements (a) and (b) of Theorem 2. The cases $k = 1$ and $k = 2$ are trivial. The cases with $3 \leq k \leq 6$ are discussed in Appendix A.

4 Split graphs and complexity results

The proofs of Theorem 3 and Theorem 4 have been moved to Appendix B and Appendix C, respectively.

References

- [1] G. AGNARSSON AND M. M. HALLDÓRSSON, *Coloring powers of planar graphs*. Proceedings of the Eleventh Annual ACM-SIAM Symposium on Discrete Algorithms (San Francisco) (2000) 654–662.
- [2] H.L. BODLAENDER, T. KLOKS, R.B. TAN, AND J. VAN LEEUWEN, λ -coloring of graphs, in Proceedings of the 17th Annual Symposium on Theoretical Aspects of Computer Science (STACS’2000), Springer LNCS 1770, (2000) 395–406.
- [3] O.V. BORODIN, H.J. BROERSMA, A. GLEBOV, AND J. VAN DEN HEUVEL, *Stars and bunches in planar graphs. Part I: Triangulations*. Preprint (2001).
- [4] O.V. BORODIN, H.J. BROERSMA, A. GLEBOV, AND J. VAN DEN HEUVEL, *Stars and bunches in planar graphs. Part II: General planar graphs and colourings*. Preprint (2001).
- [5] G.J. CHANG AND D. KUO, *The $L(2, 1)$ -labeling problem on graphs*, SIAM J. Discrete Math. 9 (1996) 309–316.
- [6] J. FIALA, A.V. FISHKIN, AND F.V. FOMIN, *Off-line and on-line distance constrained labeling of graphs*, in Proceedings of the 9th European Symposium on Algorithms (ESA’2001), Springer LNCS 2161 (2001) 464–475.
- [7] J. FIALA, T. KLOKS, AND J. KRATOCHVÍL, *Fixed-parameter complexity of λ -labelings*, Discrete Appl. Math. 113 (2001) 59–72.
- [8] J. FIALA, J. KRATOCHVÍL, AND A. PROSKUROWSKI, *Distance constrained labelings of precolored trees*, in Proceedings of the 7th Italian Conference on Theoretical Computer Science (ICTCS’2001), Springer LNCS 2202 (2001) 285–292.
- [9] D.A. FOTAKIS, S.E. NIKOLETSEAS, V.G. PAPADOPOULOU AND P.G. SPIRAKIS, *Hardness results and efficient approximations for frequency assignment problems and the radio coloring problem*, Bull. Eur. Assoc. Theor. Comput. Sci. EATCS 75 (2001) 152–180.
- [10] M.R. GAREY AND D.S. JOHNSON, *Computers and Intractability, A Guide to the Theory of NP-Completeness*, W.H. Freeman and Company, New York (1979).
- [11] M.C. GOLUMBIC, *Algorithmic Graph Theory and Perfect Graphs*, Academic Press, New York (1980).

- [12] J.R. GRIGGS AND R.K. YEH, *Labelling graphs with a condition at distance 2*, SIAM J. Discrete Math. 5 (1992) 586–595.
- [13] W.K. HALE, *Frequency assignment: Theory and applications*, Proceedings of the IEEE 68 (1980) 1497–1514.
- [14] P.L. HAMMER AND S. FÖLDES, *Split graphs*, Congressus Numerantium 19 (1977) 311–315.
- [15] J. VAN DEN HEUVEL, R.A. LEESE, AND M.A. SHEPHERD, *Graph labeling and radio channel assignment*, J. Graph Theory 29 (1998) 263–283.
- [16] J. VAN DEN HEUVEL AND S. MCGUINNESS, *Colouring the square of a planar graph*. Preprint (1999).
- [17] T.K. JONAS, *Graph coloring analogues with a condition at distance two: $L(2,1)$ -labellings and list λ -labellings*. Ph.D. Thesis, University of South Carolina (1993).
- [18] R.A. LEESE, *Radio spectrum: a raw material for the telecommunications industry*, in Progress in Industrial Mathematics at ECMI 98, Teubner, Stuttgart (1999) 382–396.
- [19] M. MOLLOY AND M.R. SALAVATIPOUR, *A bound on the chromatic number of the square of a planar graph*. Preprint (2001).
- [20] G. WEGNER, *Graphs with given diameter and a colouring problem*. Preprint, University of Dortmund (1977).

A The small cases for path backbones

Proof of the case $k=3$. Suppose that for the case $k = 3$ there is a backbone coloring of (G, T) with $\ell \leq 4$ colors. Then the equations and inequalities (3)–(6) do not have any solution s_1, s_2, s_3 over the non-negative integers. This settles the case $k = 3$.

Proof of the case $k=4$. Suppose that for the case $k = 4$ there is a backbone coloring of (G, T) with $\ell \leq 6$ colors. Then the equations and inequalities (3)–(6) have $s_1 = 3, s_2 = 0, s_3 = 1$ as unique solution over the non-negative integers. Up to symmetric cases Lemma 6.(T3) only allows $C_1 = \{1\}, C_2 = \{3\}, C_3 = \{5\}$ and $C_1 = \{1\}, C_2 = \{3\}, C_3 = \{6\}$ as mono-chromatic color sets. In the first case $C_4 = \{2, 4, 6\}$ and in the second case $C_4 = \{2, 4, 5\}$. There exists a vertex $v \in V_4$ that is adjacent to vertices from C_2 and from C_3 on the Hamiltonian path P . In either case, there is no feasible color for this vertex v , and we arrive at the desired contradiction.

Proof of the case $k=5$. Suppose for the sake of contradiction that for the case $k = 5$ there is a backbone coloring of (G, T) with $\ell \leq 7$ colors. Then the equations and inequalities (3)–(6) have $s_1 = 4, s_2 = 0, s_3 = 1$ as unique solution over the non-negative integers. By Lemma 6.(T3), the only possible mono-chromatic color sets are $C_1 = \{1\}, C_2 = \{3\}, C_3 = \{5\}, C_4 = \{7\}$. Hence, the poly-chromatic color set must be $C_5 = \{2, 4, 6\}$. But there exists a vertex $v \in V_5$ that is adjacent to vertices from C_2 and from C_3 on the Hamiltonian path P . Hence, there is no feasible color for v and we arrive at the desired contradiction.

Proof of the case $k=6$. Suppose that for the case $k = 6$ there is a backbone coloring of (G, T) with $\ell \leq 9$ colors. Then the equations and inequalities (3)–(6) have only two solutions over the non-negative integers: $s_1 = 5, s_2 = 0, s_3 = 1$, or $s_1 = 4, s_2 = 1, s_3 = 1$. Using Lemma 6.(T3), the first solution yields only one possibility for the mono-chromatic color sets, with colors 1,3,5,7,9, respectively. Since there exists a vertex v in the poly-chromatic set that is adjacent to vertices with colors 3 and 7 in P , there is no feasible color for v . We continue with the second solution. Suppose the colors c_1, c_2, c_3 and c_4 for the mono-chromatic color sets C_1, C_2, C_3, C_4 are chosen in increasing order, and let C_5 and C_6 denote the bi-chromatic and poly-chromatic color set, respectively. For a vertex $v_5 \in V_5$ and a vertex $v_6 \in V_6$ that are adjacent to vertices with colors c_2 and c_4 on P , we have no feasible color within the set $\{c_1, c_2 - 1, c_2, c_2 + 1, c_3, c_4 - 1, c_4\}$ of different colors, and we obtain an extra forbidden color if $c_4 \neq 9$. We conclude that $c_4 = 9$, and by symmetry (using c_3 and c_1) that $c_1 = 1$. If $c_3 \neq c_2 + 2$, then by considering two vertices from V_5 and V_6 that are adjacent to vertices with colors c_2 and c_3 on P , we obtain the eight forbidden colors $1, c_2 - 1, c_2, c_2 + 1, c_3 - 1, c_3, c_3 + 1$, and 9, so we cannot color both of these vertices. Hence, $c_3 = c_2 + 2$. There remain two possibilities, up to symmetry: $c_2 = 3$ (or 5) or $c_2 = 4$.

If $c_2 = 4$, we have mono-chromatic colors 1,4,6,9; we obtain a contradiction in the following way: considering vertices $v_5 \in V_5$ and $v_6 \in V_6$ adjacent to vertices with colors 1 and 6 in P , we deduce that colors 3 and 8 are not in the same set; similarly with colors 4 and 6, we deduce that colors 2 and 8 are in different sets; finally with colors 6 and 9, we obtain that colors 2 and 3 are in different sets, which is absurd.

We are left with the case that $c_2 = 3$, and with mono-chromatic colors 1,3,5,9. Using colors 3 and 5 as in the previous case, we conclude that colors 7 and 8 cannot be in the same set (V_5 or V_6); using colors 3 and 9, the same holds for colors 6 and

7; using colors 5 and 9, the same holds for colors 2 and 7. The only possibility is a bi-chromatic set $C_5 = \{4, 7\}$ and a poly-chromatic set $C_6 = \{2, 6, 8\}$. Now consider a subpath Q of P on four vertices visiting the sets in the order V_2, V_5, V_6, V_2 . Since V_2 has color 3, the only possible color on Q in V_5 is 7, and we cannot find a feasible color on Q in V_6 , our final contradiction.

B Proof of the results for Split graphs

This section is devoted to a proof of Theorem 3. The following observation is straightforward, but will be useful in many of our arguments.

Observation 7 *Let $G = (V, E)$ be a graph, let $f, g : V \rightarrow \{1, \dots, k\}$ be two colorings of V such that $f(v) + g(v) = k + 1$ for all $v \in V$. Then for any spanning tree T of G , coloring f is a backbone coloring of (G, T) if and only if g is a backbone coloring of (G, T) . ■*

Tightness of the bound in (a). Consider a split graph with a clique of k vertices v_1, \dots, v_k and with an independent set of $(k-2)(k-1)/2$ vertices $u_{i,j}$ with $1 \leq i \neq j \leq k-1$. Every vertex $u_{i,j}$ is adjacent to all vertices v_s with $s \neq i$. The spanning tree T contains the $k-1$ edges $v_k v_s$ with $1 \leq s \leq k-1$. The vertices $u_{i,j}$ form the leaves of T ; in the tree, vertex $u_{i,j}$ is adjacent only to v_j . Clearly, $\chi(G) = k$.

Suppose to the contrary that $\text{BBC}(G, T) \leq k+1$, and consider such a backbone coloring. The vertices v_1, \dots, v_k in the clique must be colored with k pairwise distinct colors. Since they form a star, either vertex v_k has color 1 and color 2 is not used on the clique, or vertex v_k has color $k+1$, and color k is not used on the clique. Both cases are symmetric as in Observation 7, and we assume without loss of generality that v_k has color $k+1$ and that color k is not used on the clique. Let v_i be the vertex that has color $k-2$, and let v_j be the vertex that has color $k-1$. The vertex $u_{i,j}$ is adjacent to all clique vertices except v_i ; hence, it could only be colored with color $k-2$ or with color k . But these two colors are forbidden for $u_{i,j}$, since in the spanning tree it is adjacent to vertex v_j with color $k-1$. Since there is no feasible color for $u_{i,j}$, we arrive at the desired contradiction.

Tightness of the bound in (b). Consider a split graph with $2k$ vertices v_1, \dots, v_{2k} . The clique C is formed by the k vertices $v_1, v_3, \dots, v_{2k-1}$ with odd index together with vertex v_{2k} . The remaining vertices form the independent set I ; every vertex in I is adjacent to all vertices in the clique C except v_{2k} . The Hamiltonian path P runs through v_1, \dots, v_{2k} by increasing index. Clearly, $\chi(G) = k+1$.

Suppose to the contrary that $\text{BBC}(G, P) \leq k+1$, and consider such a backbone coloring. Let z denote the color of v_{2k} . Let v_j denote some vertex in C that has color $z-1$ or $z+1$. Since every color is used on exactly one vertex in the clique C , every vertex in I must be colored with color z . But on the Hamiltonian path P , one of the vertices in I is adjacent to v_j , a contradiction.

Proof of the bound in (a). Let $G = (V, E)$ be a split graph with a spanning tree $T = (V, E_T)$. Let C and I be a partition of V such that C with $|C| = k$ is a clique of maximum size, and such that I is an independent set. Since split graphs are perfect, $\chi(G) = \omega(G) = k$. We consider the restriction of the tree T to the vertices in C , and we distinguish two cases.

In the first case, the restriction of T to C forms a star $K_{1, k-1}$. Let v_1, \dots, v_{k-1} denote the $k-1$ leaves of this star, and let v_k denote its center. For $i = 1, \dots, k-1$ we color v_i with color i , and we color v_k with color $k+1$. This yields a backbone

coloring for the vertices in C . All vertices $u \in I$ are leaves in the tree T . Any vertex $u \in I$ with $uv_k \notin T$ can be safely colored with color $k + 2$. It remains to consider vertices $u \in I$ with $uv_k \in T$. In the graph G , such a vertex u is nonadjacent to at least one of the vertices v_1, \dots, v_{k-1} , say to vertex v_j (otherwise, the clique C could be augmented by vertex u and would not be of maximum size as we assumed). In this case we may color u with color j .

In the second case, the restriction of T to C does not form a star. In this case the restriction of T to C has a proper 2-coloring $C = C_1 \cup C_2$ with $|C_1| = a \geq 2$ and $|C_2| = b \geq 2$. Then there exist a vertex $x \in C_1$ and a vertex $y \in C_2$ for which $xy \notin T$. Let $v_1, \dots, v_a = x$ be an enumeration of the vertices in C_1 , and let $y = v_{a+1}, \dots, v_{a+b}$ be an enumeration of the vertices in C_2 . For $i = 1, \dots, a + b$ we color vertex v_i with color i . This yields a backbone coloring of C with $a + b = k$ colors. All vertices in I are colored with color $k + 2$.

Proof of the bound in (b). Let $G = (V, E)$ be a split graph with a Hamiltonian path $P = (V, E_P)$. Let C and I be a partition of V such that C with $|C| = k$ is a clique of maximum size, and such that I is an independent set.

The case $k = 1$ is trivial. In case $k = 2$, G is a bipartite graph and $\text{BBC}(G, P) \leq 3$ by Theorem 2(a). The case $k = 4$ can be settled by (quite tedious) case distinctions. From now on we will assume that $k \geq 5$. Depending on the way the Hamiltonian path P traverses C and I in G different cases and subcases are distinguished.

Case A. There is a vertex of C with no neighbors on P in I .

If we can choose this vertex with degree two in P , let such a vertex be denoted by v ; in the other case, v denotes such an end vertex of P in C . We assign color k to v , color $k + 1$ to all vertices of I , and we claim we can extend this coloring to a backbone coloring of (G, P) using the colors $1, 2, \dots, k - 1$ for the other vertices of C . To prove this, consider the following procedure. Color the vertices of $C - \{v\}$ one by one, using the colors in decreasing order, subject to the backbone coloring restrictions, as long as this is possible. Suppose this procedure cannot be completed. Then at a certain stage color c cannot be assigned; this implies that every uncolored vertex in C is adjacent in P to the vertex with color $c + 1$. This is only possible if the number of uncolored vertices is at most two, hence $c \leq 2$. We treat these two subcases separately.

First assume $c = 2$. Then both uncolored vertices x and y are adjacent to the vertex z with color 3, and color 4 has been used on a vertex $w \notin \{x, y, z\}$ of C . At least one of the vertices x, y is not adjacent in P to w . We recolor w with color 3, assign color 4 to z , and use colors 1 and 2 for x and y , in such a way that the (possible) neighbor of w on P receives color 1.

Now assume $c = 1$. Then there is only one uncolored vertex x and it is adjacent in P to a vertex y in C with color 2. We again distinguish a number of subcases. If x has a neighbor $z \neq y$ on P in C with color $\ell \geq 4$, then we can simply recolor z with color 1, and assign color ℓ to x . In the other cases, x has a neighbor $z \neq y$ on P in C with color 3, or no neighbor on P in C . Since both colors 4 and 5 are used in C , at least one of the vertices with colors 4 and 5 is not adjacent on P to y . If we can choose such a vertex with color 5, we recolor w with color 1, and assign color 5 to x ; if we cannot choose such a vertex, then y has a neighbor u on P in C with color 5. (Then u is not adjacent to z on P in the case z is a neighbor of x on P .) We recolor u with color 2, y with color 5, and we assign color 1 to x . This completes the proof for Case A.

Case B. Every vertex of C has a neighbor on P in I .

We distinguish a number of subcases.

B1.

Let us first suppose there is no edge of P in $G[C]$, i.e. every vertex of C has all its neighbors on P in I .

If P has an end vertex v in C , we assign color k to v , color 1 to its neighbor u on P in I , and color 1 to the nonneighbor of u in C . It is easy to extend this coloring to a backbone coloring using color $k + 1$ for the remaining vertices of I , color $\neq 2$ for the uncolored neighbor of u on P in C , and the remaining colors for the other vertices of C .

If every vertex of C has degree 2 in P , we may choose such a vertex v with the property that v has two neighbors a and b on P in I , and that a and b have neighbors $v_a \neq v$ and $v_b \neq v$ on P , respectively.

If a and b have a common nonneighbor p in C , we assign color 1 to a , b , and p , color $k + 1$ to the other vertices of I , color k to v , color 3 to v_a , and color 4 to v_b . The other colors can be assigned arbitrarily to the uncolored vertices of C , yielding a backbone coloring.

We are left with the case that a and b have different nonneighbors u_a and u_b in C , respectively. We define different colorings for three situations; in all cases we assign color k to v and color $k + 1$ to all vertices of $I - \{a, b\}$.

- (i) $u_a b \notin E(P)$ and $u_b a \notin E(P)$. We assign color 1 to a , b and u_a , color 2 to u_b , and the unused colors arbitrarily to the uncolored vertices.
- (ii) $u_a b \in E(P)$ and $u_b a \in E(P)$. We assign color 1 to a and u_a , color 3 to b and u_b , and the unused colors arbitrarily to the uncolored vertices.
- (iii) $u_a b \in E(P)$ and $u_b a \notin E(P)$ (or vica versa). We assign color 1 to b and u_b , color 3 to a and u_a , color 4 to v_b , and the unused colors arbitrarily to the uncolored vertices.

This yields backbone colorings in all subcases.

B2.

In the remaining case there is a vertex v in C with one neighbor a on P in I and another neighbor on P in C . We assign color k to v , color $k + 1$ to all vertices of $I - \{a\}$, and consider the structure of the edges of P in C . These edges form a matching (of at least one edge) in $G[C]$. We extend this matching (if necessary) to a matching M of cardinality $\lfloor \frac{k}{2} \rfloor$ in $G[C]$. Now let us denote by v_a the other neighbor of a on P in C , if any, and by u_a a nonneighbor of a in C . (Possibly $u_a v \in E(P)$.)

We deal with the case $k = 5$ separately: we assign color 4 to v_a (if it exists). If u_a is adjacent to v_a (assuming it exists), we assign color 2 to u_a , and use the colors 1 and 3 for the uncolored vertices in C . If u_a is not adjacent to v_a (or if v_a does not exist), we assign color 1 to u_a , and use the unused colors for the uncolored vertices in C . In both cases we use the same color for a as for u_a , yielding a backbone coloring.

Now let us assume $k \geq 6$. We color the end vertices of the edges in M by pairs of colors $\{k - i, \lfloor \frac{k}{2} \rfloor - i\}$ for $i = 1, 2, \dots, \lfloor \frac{k}{2} \rfloor - 1$ in such a way that the colors of u_a and v_a (if it exists) differ by at least 2. It is not difficult to check that such a coloring exists. Now we assign the same color to a as to u_a , yielding a backbone coloring. This completes the proof of the final subcase.

C Proofs of the complexity results

This appendix is devoted to a proof of Theorem 4.

We start with the positive results in statement (a). So let $G = (V, E)$ be a graph with a spanning tree $T = (V, E_T)$. Let $V = V_0 \cup V_1$ be the bipartition of the vertex

set induced by T . Then in any backbone coloring with colors $\{1, 2, 3\}$, the color 2 can not be used at all. Consider some fixed vertex $v \in V_0$. By Observation 7, we may assume without loss of generality that the color of v is 1. Then all vertices in V_0 must be colored by 1, and all vertices in V_1 must be colored by 0. Hence, $\text{BBC}(G, T) = 3$ if and only if G is bipartite.

Next, consider the case of backbone colorings with $\{1, 2, 3, 4\}$. Consider some fixed vertex $v \in V_0$. By Observation 7, we may assume without loss of generality that the color of v is in $\{1, 2\}$. Then all vertices in V_0 must be colored by colors $\{1, 2\}$, and all vertices in V_1 must be colored by colors $\{3, 4\}$. Hence, $\text{BBC}(G, P) \leq 4$ if and only if the two subgraphs of G that are induced by V_0 and by V_1 are both bipartite.

Now let us prove the negative result in statement (b) of Theorem 4. The reduction is done from the NP-complete classical ℓ -coloring problem (see Garey & Johnson [10] for more information): Given a graph $H = (V_H, E_H)$, does there exist a proper ℓ -coloring of H ?

Let $H = (V_H, E_H)$ be an instance of ℓ -coloring, and let v_1, v_2, \dots, v_n be an enumeration of the vertices in V_H . We create $3(n-1)$ new vertices a_i, b_i, c_i with $1 \leq i \leq n-1$. For every $i = 1, \dots, n-1$ we introduce the new edges $v_i a_i, a_i b_i, b_i c_i$, and $c_i v_{i+1}$. The graph that results from adding these $3(n-1)$ new vertices and these $4(n-1)$ new edges to H is denoted by G . The vertices $v_1, a_1, b_1, c_1, v_2, \dots, c_{n-1}, v_n$ form a Hamiltonian path P in G . We claim that $\chi(H) \leq \ell$ if and only if $\text{BBC}(G, P) \leq \ell$.

Indeed, assume that $\text{BBC}(G, P) \leq \ell$ and consider such a backbone ℓ -coloring. Then the restriction to the vertices in V_H yields a proper ℓ -coloring of H . Next assume that $\chi(H) \leq \ell$, and consider a proper ℓ -coloring $f : V_H \rightarrow \{1, \dots, \ell\}$. We extend f to a backbone ℓ -coloring of (G, P) : Every vertex b_i receives color 3. If vertex $f(v_i) \leq 3$ then a_i is colored ℓ , and otherwise it is colored 1. If vertex $f(v_{i+1}) \leq 3$, then c_i is colored ℓ , and otherwise it is colored 1. This completes the proof of Theorem 4.