

REPORTS
IN
INFORMATICS

ISSN 0333-3590

Finding minimum feedback vertex set in
bipartite graph

Fedor V. Fomin and Artem V. Pyatkin

REPORT NO 291

February 2005



Department of Informatics
UNIVERSITY OF BERGEN
Bergen, Norway

This report has URL <http://www.ii.uib.no/publikasjoner/texrap/ps/2005-291.ps>
Reports in Informatics from Department of Informatics, University of Bergen, Norway, is
available at <http://www.ii.uib.no/publikasjoner/texrap/>.

Requests for paper copies of this report can be sent to:
Department of Informatics, University of Bergen, Høyteknologisenteret,
P.O. Box 7800, N-5020 Bergen, Norway

Finding minimum feedback vertex set in bipartite graph

Fedor V. Fomin*

*Department of Informatics, University of Bergen,
N-5020 Bergen, Norway*

fomin@ii.uib.no

Artem V. Pyatkin†

*Department of Informatics, University of Bergen,
N-5020 Bergen, Norway*

Artem.Pyatkin@ii.uib.no

17th February 2005

Abstract

We show that minimum feedback vertex set in a bipartite graph on n vertices can be found in $1.8621^n \cdot n^{\mathcal{O}(1)}$ time.

Keywords: *minimum feedback vertex set, bipartite graph, exact algorithm*

1 Introduction

Let $G = (V, E)$ be an undirected graph on n vertices. The set $X \subset V$ is called a *feedback vertex set* or an *FVS* if $G \setminus X$ is a forest. The problem of finding minimum FVS has many applications and its history can be traced back to the early 60s (see [10] for the survey). The problem is also one of the classical NP complete problem from Karp's list [12]. Thus not surprisingly, for several decades almost every new algorithmic paradigm was tried on this problem including approximation algorithms [1, 2, 9, 13], linear programming [7], local search [4], polyhedral combinatorics [6, 11], probabilistic algorithms [14], and parameterized complexity [8, 15].

In recent years the topic of exact (exponential-time) algorithms for NP hard problems has led to much research [17]. However, despite much progress on exponential-time solutions to other graph problems such as chromatic number [3, 5] or maximum independent set [16], the only worst-case bound known for finding minimum FVS is that of $\mathcal{O}^*(2^n)$ obtained by trying all possible vertex subsets. Throughout this paper we use a modified big-Oh notation that suppresses all polynomially bounded factors. For functions f and g we write $f(n) = \mathcal{O}^*(g(n))$ if $f(n) = g(n) \cdot n^{\mathcal{O}(1)}$.

In this note we present an algorithm finding a minimum FVS in a bipartite graph in $\mathcal{O}^*(1.8621^n)$ time. Note that an easy reduction (subdividing all edges) shows that the problem remains NP-hard even for bipartite graphs.

*Supported by Norges forskningsråd projects 160778/V30 and 162731/V00

†The work was partially supported by grants of the Russian Foundation for Basic Research (project codes 05-01-00395 and 05-01-00953), INTAS (project code 04-77-7173)

2 Algorithm

We need the following statement, which can be easily proved by Stirling's formula.

Proposition 2.1. *Let V be a set of cardinality n and let $0 < c \leq 1/2$. Then the number of all subsets $S \subset V$ of cardinality at most cn is bounded by $\mathcal{O}^*(1/(c^c(1-c)^{1-c})^n)$.*

First we present an algorithm finding the minimum FVS in graphs having large independent sets, and then improve it slightly for bipartite graphs.

Lemma 2.2. *Let A be an independent set in a graph G and let X be a minimum FVS in G having the largest intersection with $B = V \setminus A$. Also let $C \subseteq A$ be the set of all vertices from A having at least 3 neighbors in $B \setminus X$. Then*

- a) $X \cap A \subseteq C$;
- b) $|C| \leq \frac{3 \cdot |B \setminus X|}{2}$ and $|C \setminus X| \leq \frac{|B \setminus X|}{2}$.

Proof. Let $S = B \setminus X$ and $X_A = X \cap A$.

a) Let us note that X_A cannot contain a vertex v having at most one neighbor in S since then $X \setminus \{v\}$ would be an FVS of smaller cardinality. If a vertex $u \in X_A$ has two neighbors in S (say, v and w) then $X \cup \{v\} \setminus \{u\}$ is a minimum FVS having larger intersection with B , contradicting the choice of X . Thus $X_A \subset C$.

b) Since B is an FVS and X is a minimum FVS, we have that $|C \cap X| \leq |S|$. Let $S' = C \setminus X$. The set $S \cup S'$ induces a forest in G with at least $3 \cdot |S'|$ edges and therefore $3|S'| < |S| + |S'|$. Thus $|S'| \leq |S|/2$ and $|C| \leq 3 \cdot |S|/2$. \square

Lemma 2.3. *Let G be a graph with a given independent set of size $(1 - \varepsilon)n$. Then a minimum FVS in G can be found in $\mathcal{O}^*((1 + 3\sqrt{3}/2)^{\varepsilon n})$ time.*

Proof. Let A be an independent set of size $(1 - \varepsilon)n$ and let $B = V \setminus A$.

Consider the following procedure of finding a minimum FVS X having maximum number of vertices from B . We perform in three steps.

Step 1. First we try to guess how the set $B \setminus X$ looks like. We try all subsets $S \subseteq B$. If $G[S]$ contains a cycle then we reject this subset. Otherwise let $X_B = B \setminus S$. For every vertex $u \in A$ having exactly two neighbors (say, v and w) in S add an edge uv to $G[S]$. If the resulting graph contains a cycle we again can reject S by property a) of Lemma 2.2. If this is not the case, consider subset $C_S \subset A$ of all vertices having at least 3 neighbors in S . If $|C_S| > 3|S|/2$ we again can reject S by property b) of Lemma 2.2.

Step 2. For each set S that was not rejected at the previous step, we try all possible subsets $S' \subset C_S$ of size at most $|S|/2$. We may restrict the search to subsets of such size due to property b) of Lemma 2.2. If $S \cup S'$ induces a forest then we put $X = X_B \cup (C_S \setminus S')$ in the list of potential candidates for minimum FVS.

Step 3. Among all potential candidates we choose the one of minimum cardinality. It is a desired minimum FVS.

The correctness of the procedure follows from the fact that the list of potential candidates does not contain only the sets X which do not satisfy the conditions of Lemma 2.2.

Now let us evaluate the running time of the algorithm. At step 1 we consider all $2^{\varepsilon n}$ subsets. In the worst case, for each of them at step 2 we have to consider at most $\binom{3|S|/2}{|S|/2}$ subsets which is $\mathcal{O}^*((3/2^{2/3})^{3|S|/2})$ by Proposition 2.1. So, the overall running time of the algorithm is

$$\begin{aligned}
\sum_{s=0}^{\varepsilon n} \binom{\varepsilon n}{s} \sum_{i=0}^{s/2} \binom{3s/2}{i} &= \mathcal{O}^* \left(\sum_{s=0}^{\varepsilon n} \binom{\varepsilon n}{s} (3/2^{2/3})^{3s/2} \right) \\
&= \mathcal{O}^* \left(\sum_{s=0}^{\varepsilon n} \binom{\varepsilon n}{s} (3^{3/2}/2)^s \right) = \mathcal{O}^* \left((1 + 3\sqrt{3}/2)^{\varepsilon n} \right).
\end{aligned}$$

□

Note that $(1 + 3\sqrt{3}/2)^\varepsilon < 2$ for $\varepsilon \leq 0.54135$. Since a bipartite graph has an independent set of cardinality at least $n/2$, Lemma 2.3 yields that a minimum FVS in bipartite graphs can be found in $\mathcal{O}^*((1 + 3\sqrt{3}/2)^{n/2}) = \mathcal{O}^*(1.8968^n)$ time. In Lemma 2.3 we used only the fact that a graph contains independent set A of size $(1 - \varepsilon)n$. To prove the next theorem we exploit the fact that the graph is bipartite.

Theorem 2.4. *A minimum FVS of a bipartite graph can be found in time $\mathcal{O}^*(1.8621^n)$.*

Proof. Let $G = (V, E)$ be a bipartite graph with bipartition $V = A \cup B$, $|A| = (1 - \varepsilon)n$, $|B| = \varepsilon n$. We assume that $|B| \leq |A|$, i. e. $\varepsilon = |B|/n \leq 1/2$.

Consider the following procedure. Steps 1 and 2 of this procedure are similar to the steps from Lemma 2.3 and Steps 3 and 4 can be seen as applying the procedure from Lemma 2.3 to the set B .

Step 1. We try all subsets $S \subseteq B$ of cardinality at most $\varepsilon n/2$. Let $X_B = B \setminus S$. For every vertex $u \in A$ having exactly two neighbors (say, v and w) in S add an edge vw to S . If after that $G[S]$ contains a cycle, reject S . Otherwise, compute the subset $C_S \subseteq A$ of all vertices having at least 3 neighbors in S . If $|C_S| > 3|S|/2$ reject S .

Step 2. For each set $S \subseteq B$ that was not rejected at the first step, consider all subsets $S' \subseteq C_S$ of cardinality at most $|S|/2$. If $S \cup S'$ induces a forest then mark $X = X_B \cup (C_S \setminus S')$ as a candidate for a minimum FVS.

Step 3. Try all subsets $P' \subseteq A$. Let $Y_A = A \setminus P'$. For every vertex $u \in B$ having exactly two neighbors (say, v and w) in P' add an edge vw to P' . If after that $G[P']$ contains a cycle, reject it. Otherwise try all subsets $C'_{P'} \subseteq B$ of all vertices having at least 3 neighbors in P' . If $|C'_{P'}| > |P'|/2 + \varepsilon n/2$, reject P' .

Step 4. For each set P' that was not rejected at the third step, consider all subsets $P \subseteq C'_{P'}$ of cardinality at most $|P'|/2$. If $P \cup P'$ induces a forest then $Y = Y_A \cup (C'_{P'} \setminus P)$ is a candidate to be a minimum FVS.

Step 5. Choose among all the candidates X and Y found the one having minimum cardinality. It is a desired minimum FVS.

Let us argue first that the described procedure finds a minimum FVS. The argumentation is similar to those used in Lemma 2.3. Let X and Y be minimum FVSs having the largest intersections with B and A respectively. If $|X \cap B| \geq \varepsilon n/2$ then X is found at steps 1–2. Otherwise, $|Y \cap B| \leq |X \cap B| \leq \varepsilon n/2$, and Y will be found at steps 3–4.

We claim that the running time of the described procedure is

$$\mathcal{O}^*(\max\{(108^{\varepsilon n/4}), 2^{\varepsilon n/2}(1 + \sqrt{2})^{(1-\varepsilon)n}\}). \quad (1)$$

In fact, steps 1–2 require

$$\begin{aligned}
\sum_{s=0}^{\varepsilon n/2} \binom{\varepsilon n}{s} \sum_{i=0}^{s/2} \binom{3s/2}{i} &= \mathcal{O}^* \left(\sum_{s=0}^{\varepsilon n/2} \binom{\varepsilon n}{s} (3\sqrt{3}/2)^s \right) = \mathcal{O}^*(2^{\varepsilon n} (3\sqrt{3}/2)^{\varepsilon n/2}) \\
&= \mathcal{O}^*(108^{\varepsilon n/4}),
\end{aligned}$$

time and steps 3–4 require

$$\begin{aligned} \sum_{s=0}^{(1-\varepsilon)n} \binom{(1-\varepsilon)n}{s} \sum_{i=0}^{s/2} \binom{(\varepsilon n + s)/2}{i} &= \mathcal{O}^* \left(\sum_{s=0}^{(1-\varepsilon)n} \binom{(1-\varepsilon)n}{s} 2^{(\varepsilon n + s)/2} \right) \\ &= \mathcal{O}^* (2^{\varepsilon n/2} (1 + \sqrt{2})^{(1-\varepsilon)n}) \end{aligned}$$

time.

Now for $\varepsilon \geq 0.4856$ the theorem follows from (1) and for $\varepsilon \leq 0.4856$ the theorem follows from Lemma 2.3. \square

References

- [1] V. BAFNA, P. BERMAN, AND T. FUJITO, *A 2-approximation algorithm for the undirected feedback vertex set problem*, SIAM Journal on Discrete Mathematics, 12 (1999), pp. 289–297.
- [2] R. BAR-YEHUDA, D. GEIGER, J. NAOR, AND R. M. ROTH, *Approximation algorithms for the feedback vertex set problem with applications to constraint satisfaction and Bayesian inference*, SIAM Journal on Computing, 27 (1998), pp. 942–959.
- [3] R. BEIGEL AND D. EPPSTEIN, *3-coloring in time $O(1.3289^n)$* , Journal of Algorithms, 54 (2005), pp. 444–453.
- [4] L. BRUNETTA, F. MAFFIOLI, AND M. TRUBIAN, *Solving the feedback vertex set problem on undirected graphs*, Discrete Applied Mathematics, 101 (2000), pp. 37–51.
- [5] J. M. BYSKOV, *Enumerating maximal independent sets with applications to graph colouring*, Operations Research Letters, 32 (2004), pp. 547–556.
- [6] M.-C. CAI, X. DENG, AND W. ZANG, *A min-max theorem on feedback vertex sets*, Mathematics of Operations Research, 27 (2002), pp. 361–371.
- [7] F. A. CHUDAK, M. X. GOEMANS, D. S. HOCHBAUM, AND D. P. WILLIAMSON, *A primal-dual interpretation of two 2-approximation algorithms for the feedback vertex set problem in undirected graphs*, Operations Research Letters, 22 (1998), pp. 111–118.
- [8] R. G. DOWNEY AND M. R. FELLOWS, *Parameterized complexity*, Springer-Verlag, New York, 1999.
- [9] G. EVEN, J. NAOR, B. SCHIEBER, AND L. ZOSIN, *Approximating minimum subset feedback sets in undirected graphs with applications*, SIAM Journal on Discrete Mathematics, 13 (2000), pp. 255–267.
- [10] P. FESTA, P. M. PARDALOS, AND M. G. C. RESENDE, *Feedback set problems*, in Handbook of combinatorial optimization, Supplement Vol. A, Kluwer Acad. Publ., Dordrecht, 1999, pp. 209–258.
- [11] M. FUNKE AND G. REINELT, *A polyhedral approach to the feedback vertex set problem*, in Integer programming and combinatorial optimization, vol. 1084 of LNCS, Springer-Verlag, Berlin, 1996, pp. 445–459.
- [12] R. M. KARP, *Reducibility among combinatorial problems*, in Complexity of computer computations, Plenum Press, New York, 1972, pp. 85–103.

- [13] J. KLEINBERG AND A. KUMAR, *Wavelength conversion in optical networks*, Journal of Algorithms, 38 (2001), pp. 25–50.
- [14] P. M. PARDALOS, T. QIAN, AND M. G. C. RESENDE, *A greedy randomized adaptive search procedure for the feedback vertex set problem*, Journal of Combinatorial Optimization, 2 (1999), pp. 399–412.
- [15] V. RAMAN, S. SAURABH, AND C. R. SUBRAMANIAN, *Faster fixed parameter tractable algorithms for undirected feedback vertex set*, in Proceedings of the 13th International Symposium on Algorithms and Computation (ISAAC 2002), vol. 2518 of LNCS, Springer-Verlag, Berlin, 2002, pp. 241–248.
- [16] J. M. ROBSON, *Algorithms for maximum independent sets*, Journal of Algorithms, 7 (1986), pp. 425–440.
- [17] G. WOEGINGER, *Exact algorithms for NP-hard problems: A survey*, in Combinatorial Optimization - Eureka, you shrink!, vol. 2570 of LNCS, Springer-Verlag, Berlin, 2003, pp. 185–207.